# International **Online Conference**

# Algebraic<br>and Geometric **Methods of Analysis**

dedicate to the memory of Yuriy Trokhymchuk  $(17.03.1928 - 18.12.2019)$ 

May 25-28, 2021 Odesa, Ukraine

#### LIST OF TOPICS

- Topological methods in analysis
- Geometric problems of complex and mathematical analysis
- Algebraic methods in geometry
- $\bullet$  Differential geometry in the whole
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Geometric and topological methods in natural sciences

## ORGANIZERS

- Ministry of Education and Science of Ukraine
- Odesa National Academy of Food Technologies
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- International Geometry Center
- Kyiv Mathematical Society

#### **SCIENTIFIC COMMITTEE**

# Drozd Yu.

 $(Kyiv, Ukraine)$ Maksymenko S.  $(Kyiv, Ukraine)$ Plaksa S.  $(Kyiv, Ukraine)$ Prishlyak A.  $(Kyiv, Ukraine)$ 

# Bakhtin O.

 $(Kyiv, Ukraine)$ Balan V.  $(Bucharest, Romania)$ Banakh T.  $(Lviv, Ukraine)$ Borysenko O.  $(Kharkiv, Ukraine)$ 

Cherevko Ye.  $(Odesa, Ukraine)$ Fedchenko Yu.  $(Odesa, Ukraine)$ Karlova O.  $(Chemists, Ukraine)$ Kiosak V.  $(Odessa, Ukraine)$ Konovenko N.  $(Odessa, Ukraine)$ Lyubashenko V.  $(Kyiv, Ukraine)$ Matsumoto K.  $(Ya magata, Japan)$ Mormul P.  $(Warsaw, Poland)$ 

Mykhailyuik V.  $(Chernivtsi, Ukraine)$ Plachta L.  $(Krakov, Poland)$ Pokas S.  $(Odessa, Ukraine)$ Sabitov I.  $(Moscow, Russia)$ Savchenko O.  $(Kherson, Ukraine)$ Sergeeva A.  $(Odesa, Ukraine)$ Shelekhov A.  $(Tver, Russia)$ Zarichnyi M.  $(Lviv, Ukraine)$ 

# Special mean and total curvature of a dual surface in isotropic spaces

Ismoilov Sherzodbek Shokirjon ugli (National University of Uzbekistan, Tashkent, Uzbekistan)  $E-mail:$  sh.ismoilov@nuu.ru

In this paper, we study the properties of the total and mean curvatures of a surface and its dual image in an isotropic space. We prove the equality of the mean curvature and the second quadratic forms. The relation of the mean curvature of a surface to its dual surface is found. The superimposed space method is used to investigate the geometric characteristics of a surface relative to the normal and special normal.

Consider an affine space  $A_3$  with the coordinate system Oxyz. Let  $\overrightarrow{X}(x_1, y_1, z_1)$  and  $\overrightarrow{Y}(x_2, y_2, z_2)$ be vectors of  $A_3$ .

**Definition 1.** If the scalar product of the vectors  $\overrightarrow{X}$  and  $\overrightarrow{Y}$  is defined by the formula

$$
\begin{cases}\n(X,Y)_1 = x_1x_2 + y_1y_2 & \text{if } x_1x_2 + y_1y_2 \neq 0, \\
(X,Y)_2 = z_1z_2 & \text{if } x_1x_2 + y_1y_2 = 0,\n\end{cases}
$$
\n(1)

then  $A_3$  is said to be an isotropic space  $R_3^2$ . [1, 2]

Geometry in a plane of an isotropic space will be Euclidean if it is not parallel to the  $\alpha z$  axis. When a plane is parallel to  $oz$ , the geometry on it will be Galilean.

Since an isotropic space has an affine structure, there is an affine transformation that preserves the scalar product by formula  $(1)$ . This motion of an isotropic space is given by the formula  $[5]$ 

$$
\begin{cases}\nx' = x\cos\alpha - y\sin\alpha + a \\
y' = x\sin\alpha + y\cos\alpha + b \\
z' = Ax + By + z + c\n\end{cases}
$$
\n(2)

The second sphere is defined as a surface with the constant normal curvature. This sphere of the unit radius has the equation  $[8]$ 

$$
x^2 + y^2 = 2z,\tag{3}
$$

we call it the isotropic sphere.

Let a plane  $\pi$  be given in  $R_3^2$ , which is not parallel to the *oz* axis of the space. Consider the section of the isotropic sphere by the plane  $\pi$  and denote it by  $\Gamma$ . Since an isotropic sphere is a paraboloid of revolution, the section  $\Gamma$  by a plane is a closed curve. It was proved in [2] that  $\Gamma$  is an ellipse.

Draw tangent planes to isotropic sphere (3) through points  $M \in \Gamma$ . Denote the set of tangent planes to points F by  $\{\pi\}.$ 

The following statement holds.

**Theorem 2.** All planes of the set  $\{\pi\}$  intersect at one point. [6]

If a plane  $\pi_0$  is given by the equation

$$
z = Ax + By + C,\tag{4}
$$

then the intersection point of the planes of the set  $\{\pi\}$  will be  $(A, B, -C)$ .

**Definition 3.** The point  $(A, B, -C)$  will be called dual to plane (4) with respect to isotropic sphere  $(3)$  [6]

Let us draw the tangent plane  $\pi_M$  to the surface F at the point  $M(x_0, y_0, z_0)$ . Denote by  $M^*$  the dual image of the tangent space  $\pi_M$  with respect to the isotropic sphere. When the point  $M \in F$ changes on the surface  $F$ , its dual image describes a surface  $F^*$ .

**Definition 4.** The surface  $F^*$  is said to be the dual surface to the surface F in an isotropic space.  $[6]$ 

When F is given by the equation  $z = f(x, y)$ ,  $F^*$  has the equations

$$
\begin{cases}\nx^*(u,v) = f'_u(u,v) \\
y^*(u,v) = f'_v(u,v) \\
z^*(u,v) = u \cdot f'_u(u,v) + v \cdot f'_u(u,v) - f(u,v)\n\end{cases}
$$
\n(5)

**Lemma 5.** When the total curvature of a surface  $K = 0$ , its dual image is a point or a curve.

**Theorem 6.** The product of the total curvatures of the surface F and the dual surface  $F^*$  of the isotropic space is equal to unity:

$$
K \cdot K^* = 1. \tag{6}
$$

**Lemma 7.** The special mean curvatures of the surfaces, given by the functions  $\overrightarrow{R_1}(u, v) = f_u \cdot \overrightarrow{i} + f_v \cdot \overrightarrow{j} + f_u \cdot \overrightarrow{k}$  and  $\overrightarrow{R_2}(u, v) = f_u \cdot \overrightarrow{i} + f_v \cdot \overrightarrow{j} + f_v \cdot \overrightarrow{k}$ , are calculated, respectively, by the formulas

$$
H_m(R_1) = \frac{f_{uvv} \left(f_{uu}^2 + f_{uv}^2\right) - 2f_{uuv} \left(f_{uu} f_{uv} + f_{uv} f_{vv}\right) + f_{uuu} \left(f_{uv}^2 + f_{vv}^2\right)}{\left[f_{uu}^{\prime\prime} f_{vv}^{\prime\prime} - f_{uv}^{\prime\prime\prime}\right]^2},\tag{7}
$$

$$
H_m(R_2) = \frac{f_{vvv}\left(f_{uu}^2 + f_{uv}^2\right) - 2f_{uvv}\left(f_{uu}f_{uv} + f_{uv}f_{vv}\right) + f_{uuv}\left(f_{uv}^2 + f_{vv}^2\right)}{\left[f_{uu}''f_{vv}'' - f_{uv}''^2\right]^2}.
$$
\n(8)

**Lemma 8.** The mean curvatures of the surfaces, given by the functions  $\overrightarrow{R_1}(u, v)$  and  $\overrightarrow{R_2}(u, v)$ , are equal to zero.

**Lemma 9.** The mean curvature and special mean curvature of the dual surface (5) and the surfaces  $R_1(u, v)$ ,  $R_2(u, v)$  are connected by the equality:

$$
H_m^* = H^* + u \cdot H_m(R_1) + v \cdot H_m(R_2). \tag{9}
$$

Theorem 10. The mean curvatures defined with respect to the normal and the special normal are equal:  $H_m^* = H^*$ .

**Theorem 11.** If  $\Omega = 0$ , then the special total curvature of the surface  $F^*$  is expressed in terms of the special total curvatures of the surfaces  $F$ ,  $Z_1$ , and  $Z_2$ .

## **REFERENCES**

- [1] Artykbaev A., Sokolov, D. D. Geometry as a whole in space-time. T .: Fan. 1991.-179 p.
- [2] Artikbayev A, Ismoilov Sh. O secheniya ploskosti so izotropnogo prostranstva. Scientific Journal of Samarkand University 2020; 5(123): 84-89
- [3] Avdin M. E., Classification results on surfaces in the isotropic 3-space, AKU J. Sci. Eng., 16(2016), 239-246.
- [4] Dede, C. Ekici, and W. Goemans Surfaces of Revolution with Vanishing Curvature in Galilean 3-Space, Journal of Mathematical Physics, Analysis, Geometry 2018, Vol. 14, No. 2, pp. 141-152
- [5] Ismoilov Sh, Sultonov B. Cyclic surfaces in pseudo-euclidean space. International Journal of Statistics and Applied Mathematics 2020; 3: 28-31
- [6] Ismoilov Sh. Dual image in isotropic space. NamSU konf 2016; 1: 36-40.
- [7] Lone M.S., Karacan M.K., Dual translation surfaces in the three dimensional simply isotropic space  $I_3^1$ , Tamkang Journal of mathematics Volume 49, Number 1, 67-77, March 2018
- [8] Strubecker K., "Differentialgeometrie des isotropen Raumes II"// Math.Z.47(1942) 743-777

