

# Microbiological activity of soils with difficult ecological conditions

*Gulchekhra* Nabiyeva<sup>1\*</sup>, *Dilafuz* Makhkamova<sup>1</sup>, *Yunus* Kenjaev<sup>1</sup>

<sup>1</sup>National University of Uzbekistan Named After Mirzo Ulugbek. University str., 4, 100174 Tashkent, Uzbekistan

**Abstract.** This article presents data on irrigated meadow-alluvial soils of the Takhtakupyr region of the Karakalpak Republic. The microbiological activity (ammonifiers, spore, oligonitrophils, nitrogen fixers, actinomycetes, microscopic fungi) of meadow-alluvial soils is studied, depending on the degree of salinity. The degree of enrichment with microorganisms of highly saline, medium and slightly saline soils is determined.

## 1. Introduction

The territory of the Republic of Karakalpakstan covers the ancient, old and relatively young regions of the Amu Darya delta, and partially the Kyzyl Kum, the Ustyurt plateau and the Aral Sea. At the beginning of 2017, the area of irrigated land in the Republic of Karakalpakstan amounted to 509.6 thousand hectares (3.1%) at a total land area of 16.7 million hectares. The planted area of Karakalpakstan is mainly cultivated with cotton and grain crops. Cotton is cultivated in all southern regions of the republic, except for Muinak. Under the conditions of a wide variety of natural and geomorphological conditions, typically desert automorphic soils and azonal hydromorphic soils were formed. They consist of gray-brown, desert sandy, takyrs and takyrs, residual-hydromorphic, meadow and bog-meadow soils, as well as continental and coastal salt marshes. The study of the microbiological activity of desert soils under the influence of climate change and the degree of salinity is of great interest for soil scientists, microbiologists, ecologists and climatologists.

Biodiversity loss has become a global concern as evidence accumulates that it negatively affects the ecosystems on which human societies depend. So far, most research has focused on the environmental impacts of the loss of terrestrial biodiversity; however, much of the earth's biodiversity is literally hidden under the ground. The problem of whether the decline in biodiversity in soil communities affects the overall productivity of the ecosystem remains unresolved. It is important to investigate this considering recent observations on the reduction of soil biodiversity and changes in soil communities under intensive land use [1].

The microbiological activity of soils is a bio-indicator of soil fertility. Soil microorganisms are the key driving forces of aboveground biogeochemical cycles; however, it is still unclear how variations in the composition of the soil microbial community affect ecosystem processes [2, 3].

Hansel, C. M. et al have studied the spatial heterogeneity of the physical, chemical and biological properties of soil, which allow the spread of a variety of microbial communities [4]. Factors affecting the structuring of microbial communities, including nutrient and water availability, pH and soil structure, can vary significantly with soil depth and subsurface aggregates. They have investigated changes in microbial and functional communities in soil aggregates obtained along the soil profile.

A number of researchers refer to the correlation network analysis to study the combined presence of fungi and bacteria in soil; the differences and the coincidences in the ecosystem of these groups were found [5]. These results contradict the concept that fungi and bacteria are two separate functional groups that can be studied as isolated issues. A better understanding of the relationship between soil properties, biogeochemical cycles and interactions between fungi and bacteria is an important step towards better predicting and managing soil ecosystem functions [5].

The aim of the research is to study the number of physiological groups of microorganisms and the degree of soil

---

\*Corresponding author: [gulchekhra-nabieva@rambler.ru](mailto:gulchekhra-nabieva@rambler.ru)

enrichment with microorganisms on irrigated meadow-alluvial soils, depending on the degree of salinity.

## 2. Methods

The quantitative account for individual physiological groups of microorganisms was conducted by the generally accepted method of dilutions followed by sowing on solid elective nutrient media [6]. The accounting and study of the functional diversity of microbial communities in soils were traditionally evaluated at the level of physiological groups on the appropriate media: ammonifying bacteria on meat-peptone agar (MPA), spore bacteria on MPA with the addition of wort (1:1), oligonitrophils and nitrogen fixers on Ashby's medium, actinomycetes on starch-ammonia medium, microscopic fungi on Czapek's medium. The number of bacteria was expressed in colony-forming units per 1 g of soil (CFU/g).

## 3. Results and Discussion

The cycle of transformations of nitrogen-containing compounds in soil is inextricably linked with the development and biochemical activity of ammonifying microorganisms. Ammonifiers decompose nitrogen-containing compounds into simpler ones, which then penetrate osmotically into the cells of microorganisms and undergo deamination under the action of intracellular enzymes. All protein substances contained in soil undergo ammonification. Ammonifiers, as a rule, decompose readily available nitrogen-containing organic matter in soil. If the content of nitrogen in organic matter is less than 2%, it will be completely immobilized in the cells of microorganisms, and at a higher content, it will be assimilated with the release of ammonia [6, 7, 8, 9, 10, 11]. The ammonification process is not specific; it can be realized by groups of microorganisms belonging to different taxonomic groups: bacteria from the genera *Bacillus*, bacterium, *Pseudomonas*; fungi of the genera *Penicillium*, *Trichoderma*, *Aspergillus*; streptomycetes [12, 13]. It was found that the number of ammonifying microorganisms is greatly influenced by the modes of aeration, humidification - drying, temperature, cultivation culture, application of mineral and organic fertilizers, and especially manure [14, 15].

**Table 1.** The number of physiological groups of microorganisms in soil (CFU/g of soil)

Depth, cm	Ammonium fixers	Spore	Oligonitrophils	Nitrogen fixers	Actinomycetes	Microscopic fungi
403-section (highly saline soils)						
0-15	1800000	69000	780000	15000	30000	13000
15-30	1000000	51000	600000	-	70000	7000
30-50	800000	23000	420000	-	-	2000
2449-section (medium saline soils)						
0-15	4200000	18000	1300000	-	40000	13000
15-30	2800000	60000	1400000	-	90000	3000
30-50	2500000	7000	1000000	-	90000	3000
2448-section (highly saline soils)						
0-15	2100000	45000	790000	-	40000	3000
15-30	400000	13000	250000	40000	40000	3000
30-50	1200000	3000	480000	10000	60000	4000
404-section (slightly saline soils)						
0-15	8400000	45000	1600000	-	30000	10000
15-30	2800000	40000	1400000	-	40000	2000
30-50	1300000	15000	600000	-	40000	2000
1193-section (slightly saline soils)						
0-15	2400000	27000	1600000	60000	60000	12000
15-30	1800000	42000	1000000	40000	40000	3000
30-50	800000	9000	1000000	30000	50000	3000
1187-section (medium saline soils)						
0-15	3900000	16000	1500000	30000	30000	4000
15-30	1600000	19000	1200000	40000	70000	4000
30-50	1300000	6000	500000	30000	70000	3000

Oligonitrophilic microorganisms mineralize nitrogen-free organic matter of the soil. They are weak fixers of atmospheric nitrogen, and after the death of these microorganisms, the soil is enriched with protein nitrogen. Oligonitrophils, settling on fresh plant material, mineralize mainly the carbohydrate part of organic matter and reduce the ratio of carbon to nitrogen in the decomposed substrate [16]. The ability of oligonitrophils to develop at a very low nitrogen level provides the possibility of their development in conditions unfavorable for other microorganisms, and due to this, allows them to participate in soil processes [17]. Since, in the soil, these microorganisms almost constantly develop with an acute nitrogen deficiency, then from an environmental point of view, two features of oligonitrophils

are of greatest interest: the ability to develop at a low level of bound nitrogen in the environment and to use atmospheric nitrogen. In addition, according to Makhkamova et al [18], oligonitrophils participate in soil structuring, forming cementing mucus.


In slightly saline soils (404-1193 s), the number of ammonifiers is twice as large and ranged from 2800000 to 8400000, spore from 27000 to 45000, oligonitrophils from 10000 to 1600000, nitrogen fixers from 40000 to 60000, actinomycetes from 30000 to 40000, microscopic fungi from 10000 to 12000 CFU/g of soil. In medium saline soils (2449-1187 s), the number of ammonifiers ranged from 2500000 to 4200000, spore from 7000 to 18000, oligonitrophils from 18000 to 1300000, actinomycetes from 40000 to 90000, microscopic fungi from 3000 to 13000 CFU/g of soils. In highly saline soils (403-2448 s), the number of ammonifiers ranged from 1800000 to 2100000, spore from

69000 to 45000, oligonitrophiles from 780000 to 790000, nitrogen fixers from 4000 to 15000, actinomycetes from 30000 to 40000, microscopic fungi from 3000 to 13000 CFU/g of soils. The number of microorganisms from the upper horizons to the lower horizons decreased gradually (Table 1).

The results of the study showed that in highly saline soils, bacteria-ammonifiers, according to the level of enrichment by the Zvyagintsev scale [19], are depleted in 0-15, 15-30 cm layers and very depleted in 30-50 cm. In medium saline soils in 0-15 cm layer, the number of bacteria is at an average level, and the layers of 15-30 cm and 30-50 cm are depleted and very depleted, respectively. In slightly saline soils in the 0-15 cm layer, the number of bacteria is at an average level, and 15-30 cm and 30-50 cm layers are depleted.


In highly saline and slightly saline soils, actinomycetes, according to the level of enrichment by the Zvyagintsev scale, in the layers of 0–15, 15–30, 30–50 cm, the number of bacteria is at an average level. According to the results of the study, microscopic fungi (by the Zvyagintsev scale) in highly saline soils in 0-15, 15-30, 30-50 cm layers are depleted, in medium saline soils in 0-15, 15-30 cm layers, the number of bacteria is at an average level, in 30-50 cm layer it is depleted. Slightly saline soils in 0-15 cm layer are medium supplied, in 15-30, 30-50 cm layers they are depleted.

The formation and development of soils in the desert region are characterized by constant variability associated with environmental conditions. The process of salinization of these soils affects not only the properties of soil but also the flora; this leads to a decrease in microbiological activity. We know that the study of microflora in soils is an important diagnostic index of soils (Figure 1).



**Fig. 1.** The number of physiological groups of microorganisms in soil: Ammonium fixers, Spore, and Oligonitrophils

The study revealed the presence of agronomically important groups of microorganisms, including *ammonifiers*, *spores*, *actinomycetes*, *microscopic fungi* in the composition of irrigated soils in Bukhara district of Bukhara region (Figure 2).



**Fig. 2.** Groups of microorganisms in irrigated soils of the experimental field (Note: A. *Ammonifiers*; B. *Spores*; B. *Colonies of actinomycetes*.)

According to the diagnostic indices obtained, from an environmental point of view, soil microbiology is severely damaged, which negatively affects the entire process in soil, leading to a long-term decrease in soil fertility and the loss of generations of microorganisms, as well as to a complete loss of fertility in some areas.

#### 4. Conclusion

The number of ammonifiers (8400000- 4200000) and spore (45000 -18000) in slightly saline soils is twice as high as in medium saline soils. The number of oligonitrophiles in slightly saline soils is 1600000, in medium saline soils, it is 1300000; the lower number of nitrogen fixers is explained by the lack of nutrients and extreme conditions in the area under study. The number of actinomycetes in slightly saline soils is 30000; in medium saline soils, it is  $4 \times 10^4$ , microscopic fungi in slightly saline soils is 10000; in medium saline soils, it is 13000 CFU/g of soil.

In highly saline soils, ammonifying bacteria are depleted in terms of enrichment by the Zvyagintsev scale, medium saline soils are moderately supplied, and slightly saline soils are enriched. In highly saline, medium and slightly saline soils, actinomycetes in terms of enrichment by the Zvyagintsev scale turned out to be moderately supplied, while microscopic fungi in highly saline soils are depleted; slightly saline soils are moderately supplied, and in 15-30, 30-50 cm layers they are depleted.

#### References

1. Wagg C, Bender SF, Widmer F, Van Der Heijden MGA, Soil biodiversity and soil community composition determine ecosystem multifunctionality, *National Academy of Sciences of the United States of America* **111**(14), 5266-5270 (2014)
2. Leff JW, Nemergut DR, Grandy AS, O'Neill SP, Wickings K, Townsend AR, Cleveland CC, The Effects of Soil Bacterial Community Structure on Decomposition in a Tropical Rain Forest, *Ecosystems* **15**(2), 284-298 (2012)
3. Sharipov O, Maxkamova D, Gafurova L, Hydromorphic soils of the desert zone and the biological basis for increasing their fertility, Biodiversity: global and regional processes, *Proceedings of the All-Russian Conference of Young Scientists with International Participation. Institute of General and Experimental Biology SO RAS, Moscow* (2016)
4. Hansel CM, Fendorf S, Jardine PM, Francis CA, Changes in bacterial and archaeal community structure and functional diversity along a geochemically variable soil profile, *Applied and Environmental Microbiology* **74**(5), (1620-1633) 2008.
5. de Menezes AB, Richardson AE, Thrall PH, Linking fungal-bacterial co-occurrences to soil ecosystem function, *Current Opinion in Microbiology* **37**, 135-141 (2017)
6. Bab'eva II, Zenova GM, Biology of soils, *Moscow State University, Moscow* (1989)
7. Atoev B, Kaypnazorov J, Egamberdieva M, Makhhammadiev S, Karimov M, Makhkamova D, Technology of nutriating winter wheat varieties in variety-soil-fertilizer system, *E3S Web of Conferences* **244**, 02040 (2021)

8. Makhkamova D, Gafurova L, Nabieva G, Makhhammadiev S, Kasimov U, Juliev M, Integral indicators of the ecological and biological state of soils in Jizzakh steppe, *IOP Conf. Series: Earth and Environmental Science* **1068**, 012019 (2022)
9. Gafurova LA, Madrimov RM, Razakov AM, Nabiyeva GM, Makhkamova DYu, Matkarimov TR, Evolution, Transformation and Biological Activity of Degraded Soils, *Int J Advanced Science and Technology* **28**(14), 88-99 (2019)
10. Gafurova LA, Maxkamova DYu, Gypsum-bearing soils of the Jizzakh steppe and their biological activity, *RGAU-MSXA* **34**, 76 (2019)
11. Maxkamova DYu, Seasonal variation of ammonifier bacteria in heavy meliorated soils, *Innovation Technical and Technology* **2**(1), 54-58 (2021)
12. Andryuk EI, Iutinskaya GA, Dulgerov AN, Soil microorganisms and intensive land use, *Naukova Dumka*, Kiev (1988)
13. Nabieva GM, Botirova NT, Microbiological activity of irrigated meadow-alluvial soils of Takhtakupirsky region, *Scientific integration in the interpretation of the modern educational process of the third millennium*, Kazan (2019)
14. Devyatova TA, Bezler NB, Antonyuk AN, The influence of agricultural land and soil subtype on the microbial community of zonal soils on the example of the black earth of the stone steppe, *Vestnik VSU. Series of Chemistry, Biology, Pharmacy* **1**, 46-49 (2003)
15. Maxkamova D, Gafurova L, The influence of soil salinization and gypsum on the activity of microorganisms in the soils of the Golodnaya desert, Biodiversity: global and regional processes, *All-Russian conference of young scientists with international participation. Institute of General and Experimental Biology SO RAS*, Moscow (2016)
16. Trypolskaya LN, Bagdanavichene ZE, Romanovskaya DK, Microbiological activity of sod-podzolic soil and decomposition of organic fertilizers in the autumn-winter period, *Pochvovedenie* **9**, 1100-1108 (2004)
17. Germanova NI, Medvedeva MV, Microflora of soils of the "Kivach" reserve, *Proceedings of the Karelian Scientific Center of the Russian Academy of Sciences* **10**, 10-33 (2006)
18. Makhkamova D, Nabiyeva G, Abdushukurova Z, Iskhakova Sh, Abdujabbarovna A, Climate conditions, hydrogeology and meliorative conditions of serozem -grass soils of Mirzaabad district, Sirdaryo region, *E3S Web of Conf.* **413**, 03033 (2023)
19. Zvyagintsev DG, Biological activity of soils and scales for assessing some of its indicators, *Pochvovedenie* **6**, 48-54 (1978)